



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	Books
Search	PubMed	<input type="checkbox"/> for					Go	Clear
		Limits	Preview/Index	History	Clipboard	Details		

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

Entrez
PubMed☐ 1: Allergy 2002 Apr;57(4):323-8

Related Articles, Books, LinkOut

**Down-regulated IL-5 receptor expression on peripheral blood eosinophils from budesonide-treated children with asthma.****Hellman C, Lonnkvist K, Hedlin G, Hallden G, Lundahl J.**PubMed
ServicesDepartment of Medicine, Division of Clinical Immunology and Allergy,
Stockholm, Sweden.Related
Resources

BACKGROUND: The expression and function of cytokine receptors on peripheral blood eosinophils (PBE) from healthy and asthmatic children are poorly characterized. **METHODS:** The PBE count and expression of IL-5 receptor (R) and GM-CSFR positive PBE was analyzed in nonsteroid-treated asthmatic children (n = 13), budesonide-treated asthmatic children (n = 24) and healthy children (n = 16) by flow cytometry. Alterations in intracellular EG2-epitope expression were used to measure the in vitro responsiveness of PBE to recombinant IL-5 and GM-CSF. **RESULTS:** The PBE count was increased ($P < 0.05$) in both asthmatic groups, independent of treatment, as compared to healthy children. The IL-5R expression on PBE, as well as the in vitro responsiveness of PBE to recombinant IL-5, was reduced ($P < 0.05$), in budesonide-treated asthmatic children compared to nonsteroid-treated asthmatic children and healthy children. The proportion of GM-CSFR positive PBE and in vitro responsiveness of PBE to recombinant GM-CSF were not different between the groups. In vitro treatment with budesonide did not down-regulate the proportion of IL-5R positive PBE. **CONCLUSIONS:** Budesonide-treatment of asthmatic children induces a selectively reduced IL-5R expression on PBE, concomitant with a reduced in vitro responsiveness of PBE to IL-5. We suggest that this budesonide-related down-regulation of the IL-5R might be a mechanism by which steroid treatment inhibits the action of IL-5 on eosinophil accumulation and activation in vivo.

PMID: 11906363 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="checkbox"/>	Sort	<input type="checkbox"/>	Save	Text	Clip Add	Order
---------	----------	--------------------------	------	--------------------------	------	------	----------	-------

Write to the Help Desk